

## Freezer Safety

A full freezer will remain below 40°F for about two days, and a less-than-full freezer will maintain that temperature for approximately one day. Rearrange meats so their juices do not drip onto other foods as the meats begin to thaw. Throw away any ready-to-eat item that comes in contact with meat juices.




Frozen foods that have partially or completely thawed can be refrozen if they contain ice crystals. If they have completely thawed but are still at a temperature of 40°F or below, they also can be refrozen. Use a digital thermometer to check the temperature of the food. Although partial thawing and refreezing of these foods will be safe, the quality of some foods, especially vegetables and fruits, may be reduced. Hard cheeses, breads, and fruits and vegetables and their juices—if they

look and smell normal—can be refrozen even if they have been above 40°F for more than two hours.

One way to ensure the safety of thawed meat products that have been above 40°F for two hours or less is to cook them immediately. Either serve the food immediately or refreeze the cooked item. Be sure to cook to the proper temperatures, checking internal temperatures with a digital thermometer. Refrigerate and use within two days or freeze the leftovers immediately.

Foodborne illness is very uncomfortable and sometimes life-threatening, especially for the very young, the senior population, and those with weakened immune systems. Don't take chances on your family's safety for the sake of a few dollars in groceries. Remember: When in doubt, throw it out!



## For more food safety information, contact

Penn State Cooperative Extension on the Web at [www.extension.psu.edu](http://www.extension.psu.edu)

Prepared by Mary Alice Gettings, nutrition and food safety educator, Penn State Cooperative Extension in Beaver County.

## Resources

Penn State Food Safety Web Site  
[foodsafety.cas.psu.edu](http://foodsafety.cas.psu.edu)

USDA Food Safety and Inspection Service  
U.S. Department of Agriculture  
Washington, D.C. 20250-3700  
[www.fsis.usda.gov](http://www.fsis.usda.gov)

Meat and Poultry Food Safety Hotline  
Phone: 1-800-535-4555

Visit Penn State's College of Agricultural Sciences on the Web:  
[www.cas.psu.edu](http://www.cas.psu.edu)

Penn State College of Agricultural Sciences research, extension, and resident education programs are funded in part by Pennsylvania counties, the Commonwealth of Pennsylvania, and the U.S. Department of Agriculture.

This publication is available from the Publications Distribution Center, The Pennsylvania State University, 112 Agricultural Administration Building, University Park, PA 16802. For information telephone 814-865-6713.

This publication is available in alternative media on request.

The Pennsylvania State University is committed to the policy that all persons shall have equal access to programs, facilities, admission, and employment without regard to personal characteristics not related to ability, performance, or qualifications as determined by University policy or by state or federal authorities. It is the policy of the University to maintain an academic and work environment free of discrimination, including harassment. The Pennsylvania State University prohibits discrimination and harassment against any person because of age, ancestry, color, disability or handicap, national origin, race, religious creed, sex, sexual orientation, gender identity, or veteran status. Discrimination or harassment against faculty, staff, or students will not be tolerated at The Pennsylvania State University. Direct all inquiries regarding the nondiscrimination policy to the Affirmative Action Director, The Pennsylvania State University, 328 Boucke Building, University Park, PA 16802-5901; Tel 814-865-4700/V, 814-863-1150/TTY.

Produced by Ag Communications and Marketing

© The Pennsylvania State University 2008

Code # **UK065** Rev10M10/08mpc4346

# Keeping Food Safe When the Power Goes Out!



PENNSTATE



College of Agricultural Sciences  
Agricultural Research and  
Cooperative Extension

As a result of thunderstorms, ice storms, and other disturbances, power outages occur more often than we desire. Outages are not only an inconvenience, but also a safety issue, particularly when it comes to food. When refrigerators and freezers lose power, we worry about whether or not the food will be safe for our families. Knowing how to handle food while the power is off and when it has been restored will help to ensure that you and your family do not become victims of a foodborne illness.

To monitor temperatures, it's important to keep an appliance thermometer in both the refrigerator and the freezer at all times. This will take the guesswork out of determining the temperature of your units. Also, keep a digital thermometer on hand. This utensil will enable you to check the temperature of thawed foods to determine their safety.

When the power goes out, try to find out how long it is expected to be off. If it will be for only a few hours, don't do anything. Open the freezer or refrigerator door as little as possible, for this will allow the units to maintain their chilled temperatures longer.



## Refrigerator Safety

If the power is off for more than two hours, you should begin to monitor the temperature of the refrigerator. Keep the thermometer close to the opening of the door. This will allow you to open the door only slightly, keeping the temperature lower. Check the temperature each hour. When it reaches 40°F, place block ice in a container in the refrigerator. You also might consider placing perishable items in a cooler with ice or ice packs around them.

If the temperature of the refrigerator rises above 40°F for more than two hours, you will need to discard the perishable foods listed below:

- meat, poultry, seafood
- cold cuts, hot dogs
- casseroles
- soft cheeses, shredded cheeses, low-fat cheeses
- cream, sour cream
- yogurt, milk
- eggs
- custards, pudding
- creamy salad dressings, fish sauces, hoisin sauce, opened spaghetti sauce
- cookie dough, refrigerator biscuits or rolls
- cooked pasta, rice, pasta salads
- pizza
- cut fresh fruits
- greens, cooked vegetables, opened vegetable juice
- baked or mashed potatoes, potato salad

Any food that has an unusual odor, color, or texture also should be discarded. Never taste foods to determine their safety!

Foods that are safe, even if held at 40°F or above for more than two hours, are:

- hard and processed cheeses that are properly wrapped
- grated Parmesan and Romano cheeses
- margarine that is properly wrapped
- canned or uncut fresh fruit and fruit juices (but discard if they look, feel, smell, or taste unusual)
- fresh vegetables, canned vegetables
- herbs, spices
- peanut butter, jelly, relish, mustard, ketchup, olives
- taco, barbecue, and soy sauces
- breads, rolls, cakes, muffins, bagels, pancakes, waffles
- opened jars of mayonnaise, tartar sauce, or horseradish (unless held at 50°F for more than eight hours—then throw them away)

**Bacteria grow between the temperatures of 40°F and 140°F.**

